



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

June 2, 2015

Lisa Adamson  
Regulatory Group Leader  
Control Solutions Inc.  
5903 Genoa Red Bluff  
Pasadena, Tx 77507-1041

Subject: Notification per PRN 98-10 – Add State specific language and marketing claims  
Product Name: PRODIAMINE 0.37 PLUS  
EPA Registration Number: 53883-172  
Application Date: 3/23/15  
Decision Number: 505014

Dear Ms. Adamson:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “Notification” and will be placed in our records.

If you have any questions, you may contact Dominic Schuler at 703-347-0260 or via email at [schuler.dominic@epa.gov](mailto:schuler.dominic@epa.gov).

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague". The signature is written in a cursive style with a long horizontal line extending from the end of the name.

Kathryn Montague, Product Manager 23  
Herbicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

## PRODIAMINE 0.37 PLUS

[with Lawn Fertilizer]

*Optional Marketing Language:*

[FOR PRE-EMERGENCE CONTROL OF GRASS AND BROADLEAF WEEDS IN LAWNS AND ORNAMENTAL TURF(excluding golf course putting greens)]

[LANDSCAPE ORNAMENTALS, ESTABLISHED PERENNIALS AND WILDFLOWER PLANTINGS]

[STOPS MANY WEED GRASSES BEFORE THEY GROW  
PREVENTS CRABGRASS, ANNUAL BLUEGRASS, FOXTAIL]

[Prevents Crabgrass in Lawns]  
[Phosphorus-Free][Helping our water sources]  
[With Lawn Fertilizer]  
[Controls crabgrass and broadleaf weeds before they germinate]  
[Feeds for 3 Months guaranteed!]  
[Prevent crabgrass and fertilize your lawn in one application]  
[Up to 5 month weed and grass control]  
[Covers 5,000 sq. ft.] for 15 lb. bag  
[Covers 15,000 sq ft] for 45 lb. bag  
[Covers 17,500 Square Feet] for 50 lb bag  
Prevents up to 200 Weeds  
Instantly activated by rain or irrigation  
Apply in Early Spring & Late Fall  
Plus Lawn Food  
Quick Spring Green Up after Winter  
Stops Crabgrass before it starts  
Fast Acting Formula  
Prevents Crabgrass All Season Long  
Inhibits weed seeds germination and root development  
Feeds up to 4 months guaranteed  
Phosphorus Free  
Up to 4 Month Feeding  
Controlled-Release Fertilizer  
Driven By Duration CR  
Engineered for Performance  
For a beautiful [Product name] lawn

### NOTIFICATION

53883-172

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

06/02/2015

---

#### ANNUAL FEEDING PROGRAM:

IMPORTANCE OF FEED: Regularly feeding your lawn provides the necessary nutrients it needs to help fight weeds, heat, drought and traffic.

EARLY SPRING FEEDING: After the first time you mow, feed to help your lawn green-up quickly after the harsh winter months.

LATE SPRING FEEDING: Before the hot summer months, feed to prepare and strengthen for stresses of heat, drought and traffic.

EARLY FALL FEEDING: After your lawn comes out of dormancy, feed for a quick recovery from summer stress.

LATE FALL FEEDING: Once the leaves have fallen, feed to build a strong root system and prepare your lawn for winter.

**ACTIVE INGREDIENT:**

Prodiamine: (CAS # 29091-21-2).....0.37%

**OTHER INGREDIENTS:** ..... 99.63%

TOTAL: ..... 100.00%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See back panel for further precautions and First Aid information

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

EPA REG NO. 53883-172

EPA EST. NO.

Control Solutions, Inc.  
5903 Genoa-Red Bluff Rd.  
Pasadena, TX 77507-1041  
281-892-2500

NET CONTENT: \_\_\_\_\_

<b>FIRST AID</b>	
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>● Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>● Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>● Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>● Take off contaminated clothing.</li> <li>● Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>● Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>● Call a poison control center or physician immediately for treatment advice.</li> <li>● Have person sip a glass of water if able to swallow.</li> <li>● Do not induce vomiting unless told to do so by the poison control center or physician.</li> <li>● Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>● Move person to fresh air.</li> </ul>

	<ul style="list-style-type: none"><li>● If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li><li>● Call a poison control center or doctor for further treatment advice.</li></ul>
<p style="text-align: center;"><b>HOT LINE NUMBER</b></p> <p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® International (866) 897-8050 for emergency medical treatment information.</p>	

### ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

### DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This [XXXX] lb bag covers [XXXX] sq ft of lawn area. A lawn typically will utilize 1 to 4 lbs of nitrogen per 1,000 sq ft per growing season. When used as directed, this product will supplement your regular fertilizer program by providing [XXXX] lb of nitrogen per [XXXX] sq ft. Please consult your local extension office for questions regarding a regular fertilizer program.

Apply only to actively growing turf. Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands in the driveway, sidewalk, or street, back onto your lawn. Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.

This product is a pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds, germination and root development. When used according to directions, it will control; barnyardgrass; bluegrass; annual (*Poa annua*); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass<sup>1</sup>; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass<sup>3</sup>, shepherd's purse<sup>2</sup>, signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).


<sup>1</sup>In many areas a single application of 4.1 to 9.3 pounds/1,000 square feet (175 to 405 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 4.1 to 6.2 pounds/1,000 square feet (175 to 270 pounds/acre), followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).

<sup>2</sup>For this weed, apply in late summer, fall or winter prior to germination.

<sup>3</sup>Suppression only.

**WHEN TO APPLY TO ESTABLISHED TURF:** This product must be applied prior to germination of weed seeds, as it will not control established weeds.

**THIS MAP SHOWS APPROXIMATE CRABGRASS GERMINATION DATES**



**APPLICATION DIRECTIONS:** Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days of application.

**USE PRECAUTIONS:** The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil).

Do not apply newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

**RATES OF APPLICATION:** This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

<b>MINIMUM AND MAXIMUM APPLICATION RATES (by Turf Grass Species)</b>		
<b>Turf Species</b>	<b>Minimum Rate</b>	<b>Maximum Annual Rate<sup>1</sup></b>
Bahiagrass, Bermudagrass <sup>2</sup> Centipedegrass, Zoysiagrass, Seashore Paspalum, St. Augustinegrass Tall Fescue (including turf- Type)	2.7 lbs/1,000 sq. ft. (118 lbs/acre)	9.3 lbs/1,000 sq. ft. per year (405 lbs/acre per year)
Buffalograss Kentucky Bluegrass Perennial Ryegrass	2.7 lbs/1,000 sq. ft. (118 lbs/acre)	6.2 lbs/1,000 sq. ft. per year (270 lbs/acre per year)
Creeping Red Fescue	2.7 lbs/1,000 sq. ft. (118 lbs/acre)	4.6 lbs/1,000 sq. ft. per year (202 lbs/acre per year)
Creeping Bentgrass	2.7 lbs/1,000 sq. ft. (118 lbs/acre)	4.0 lbs/1,000 sq. ft. per year (175 lbs/acre per year)

<sup>1</sup>These are the maximum rates per calendar year.

<sup>2</sup>May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 3.3 pounds/1,000 square feet (145 pounds/acre). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

**WHEN TO OVERSEED AFTER APPLICATION:** This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/reseeding results. (See map above.)

<b>USE RATE</b>	<b>MONTHS BEFORE OVERSEEDING</b>		
	<b>North</b>	<b>Transition</b>	<b>South</b>
3 pounds/1,000 sq. ft. (135 pounds/acre)	4	4	4
4 pounds/1,000 sq. ft. (175 pounds/acre)	5	4	4
4.6 pounds/1,000 sq. ft. (202 pounds/acre)	6	5	5
4.9 pounds/1,000 sq. ft. (215 pounds/acre)	-	6	6
6.2 pounds/1,000 sq. ft. (270 pounds/acre)	-	7	7
7 pounds/1,000 sq. ft. (308 pounds/acre)	-	-	9
8 pounds/1,000 sq. ft. (350 pounds/acre)	-	-	10
9.3 pounds/1,000 sq. ft. (405 pounds/acre)	-	-	12

**LANDSCAPE ORNAMENTALS:** (including established perennials and wildflower plantings): This product may be applied for residual pre-emergence weed control in ornamentals. Apply at the rate of 2.7 to 9.3 pounds/1,000 square feet (118 to 405 pounds/acre) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 9.3 pounds/1,000 square feet (405 pounds/acre) per year.

<b>SPREADER SETTINGS</b> to apply at the listed pounds per 1,000 square feet						
	3 lbs.	4 lbs.	4.6 lbs.	6.2 lbs.	8 lbs.	9.3 lbs.
Spreader A	X	X	X	X	X	X
(May contain additional spreader brands/models, spreader settings and spreader use instructions as appropriate for the product)						

**APPLICATION TIMING AND INFORMATION:** This product may be applied to newly transplanted and established ornamentals as a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate post-emergence herbicide.

**TOLERANT ORNAMENTAL SPECIES:** This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

**SCIENTIFIC NAME**

Abies spp.  
Abelia grandiflora  
Acer palmatum  
A. Platanoides  
Achillea spp.  
Actinidia chinensis  
Agapanthus africanus  
Agapanthus orientalis  
Akebia quintata  
Allium cemuum  
Anemone hybrida  
Aquilegia spp.  
Arctostaphylos densiflora  
Arctotheca calendula  
Artemesia spp.  
Aster spp.  
Aster X Frikartii  
Athrium Filiz-femina  
Aucuba japonica  
Begonia spp.  
Berberis gladwynensis  
B. Julianae  
B. mentorensis  
B. Thunbergii  
B. Verruculosa  
Bergenia cordifolia  
Boltonia asteroides  
Buddleia davidii  
Buxus microphylla  
Callistemon viminalis  
Calluna vulgaris  
Campanula carpatica  
Campis X Tagliabuana  
Carpobrotus edulis  
Cassia artemisoides

**COMMON NAME**

Fir species\*\* (Balsam, Frasier, Noble, etc.)  
Abelia: Sherwood  
Japanese Maple  
Norway Maple  
Yarrow: King Edward  
Kiwi\*  
Lily-of-the-Nile (African Lily)  
  
Five-leaf or Chocolate Vine  
Lady's Leek, Nodding Onion  
Japanese Anemone  
Aquilegia: Red and Gold  
Vine Hill Manzanita  
Cape Weed  
Wormwood; Silver Mound, Castle  
Aster: Bonnie Blue, Purple Dome  
  
Lady Fern; Fern Lady  
Japanese Aucuba  
Fibrous Begonia: Hardy Grandis  
Barberry  
Wintergreen Barberry  
Mentor Barberry  
Japanese Barberry  
Warty Barberry  
  
Snowbank  
Butterfly-Bush (Dwarf Blue); Royal Red  
Japanese Boxwood  
Weeping Bottlebrush  
Scotch Heather  
Tussock Bellflower, (White Clips)  
Trumpet Flower, Madame Galen  
Hottentot Fig (Ice Plant)  
Feathery Cassia

Ceanothus rigidus	Wild Lilac
Ceratostigma plumbagonoides	
Chamaecyparis pisifera	False Cypress
Cleyera japonica	Cleyera
Citrus spp.	Citrus species*
Coreopsis spp.	Coreopsis (Calliopsis) Early Sunrise, Moonbeam
Cornus florida	Flowering Dogwood
C. Stolonifera	American Dogwood
Cortaderia selloana	Pampas Grass
Cotoneaster apiculatus	Cranberry Cotoneaster
C. buxifolius	Cotoneaster
C. dammeri	Bearberry Cotoneaster
C. Microphyllus	Rockspray Cotoneaster
Cretaegus spp.	Hawthorne
Crocosmia spp.	Lucifer
Cupressus sempervirens	Italian Cypress
Delosperma alba	White Trailing Ice Plant
Delosperma spp.	Cooperi Pink
Delphinium spp.	Larkspur: Blue Elf
Dianthus deltoides	Dianthus: Maiden Pinks “Zing”
D. gratianopolitanus	Cheddar Pink
Dodonea viscosa	Hop Bush
Echinacea purpurea	Coneflower, Purple; Magnus
Elaeagnus pungens	Silverberry
Euonymus fortunei	Wintercreeper
E. japonica	Evergreen Euonymus
E. kiautschovica	Spreading Euonymus
Fatsia japonica	Japanese Aralia
Forsythia intermedia	Border Forsythia
Forsythia suspensa	Weeping Forsythia
F. viridissima	Greenstem Forsythia
Gaillardia spp.	Gaillardia, Blanket Flower: “Goblin”
Gardenia jasminoides	Gardenia, Cape-Jasmine
Gaura spp.	
Gentiana dahurica	Gentian
Geranium cinereum	Cranesbill
Gladiolus spp.	Gladiolus species**
Gypsophilia repens	Baby’s Breath
Hedera helix	English Ivy
Hellanthemum spp.	Sunrose
Hemerocallis spp.	Daylily: Aztec Gold, Stella de Oro, Tender Love
Hibiscus	Rose of Sharon**
Hibiscus spp.	Mallow: Disco Belle White
Hibiscus rosa-sinensis	Chinese Hibiscus
Hosta plantaginea	Hosta, Plantain Lily (Fragrant)
H. Sieboldiana	Hosta, “Seersucker”
Houttuynia cordata var. variegata	
Hydrangea macrophylla	Bigleaf Hydrangea
Ilex Bernyi	Holly
Ilex cornuta	Chinese Holly
I. crenata	Japanese Holly
I. opaca	American Holly
I. vomitoria	Yaupon Holly, Schillings
Inula ensifolia	
Iris ensata	Sword-Leaved Iris; Jodlesong
Iris siberica	Siberian Iris; Cabernet

4-20-2015 Notification to EPA – Add state restrictive language and optional marketing claims

Iris spp.	Iris species**
Jasminium nudiflorum	Winter Jasmine
Juniperus chinensis	Chinese Juniper; Nick's Compact, Parsonii
J. conferta	Shore Juniper; Blue Pacific
J. davurica	Parsoni
J. horizontalis	Creeping Juniper
Juglans spp.	Walnut*
Justicia brandegeana	Shrimp Plant
Lagerstromia indica	Crape Myrtle
Lagerstromia indica X Fauriei	Crape Myrtle; Tuscarora
Lantana montevidensis	Weeping Lantana
Lavender spp.	Lavender; Munstead
Leonopodium alpinum	Edelweiss
Ligustrum amurense	Amur Privet
Ligustrum sinense	Chinese Privet; Variegata
L. japonicum	Japanese Privet
L. lucidum	Glossy Privet (Waxleaf)
Lilium spp.	Lily; Jazz
Liriope muscari	Big Blue Lily, Silver Mound, Evergreen Giant
Liriope muscari var. variegata	Liriope, Variegated
Liriope spicata	Liriope, Creeping
Lobelia Cardinalis	Cardinal Flower, Indian Pink
Lonicera japonica	Japanese Honeysuckle
L. tatarica	Tatarian Honeysuckle
Lorapetalum chinense	Burgundy
Lythrum spp.	Loosestrife; Modern Pink
Magnolia spp.	Magnolia species**
Maleophora luteola	Ice Plant
Malus spp.	Crabapple*
Miscanthus sinensis	Yaku Jima, Silberfeder
Nandina domestica	Heavenly Bamboo
Narcissus spp.	Narcissus species**
Nerium oleander	Oleander
Olea europaea	Olive*
Ophiopogon japonicus	Mondo Grass
Omanthus heterophyllus	Osmanthus (False Holly); Gulf Tide
Osteospermum fruticosum	Trailing African Daisy
Oxydendrum arboretum	Sourwood
Paeonia suffruticosa	Tree Peony
Pennisetum setaceum	Fountain Grass (Dwarf)
Perovskia atriplicifolia	
Persea americana	Avocado*
Photinia Fraseri	Frasier's Photinia (Redtip)
Physotegia Virginiana	Dragonhead, False; Vivid
Picea spp.	Spruce species** (Colorado Blue, Norway, etc.)
Pieris japonica	Lily-of-the-Valley Bush
Pinus brutia	Calabrian Pine
P. canariensis	Canary Island Pine
P. Elliottii	Slash Pine
P. halepensis	Aleppo Pine
P. nigra	Austrian Pine
P. palustris	Longleaf Pine
P. radiata	Monterey Pine
P. strobus	Eastern White Pine
P. sylvestris	Scotch Pine
P, taeda	Loblolly Pine

4-20-2015 Notification to EPA – Add state restrictive language and optional marketing claims

P. Thunbergiana	Japanese Black Pine
P. Virginiana	Virginia Pine
Pistachio spp.	Pistachio*
Pittosporum rhombifolium	Queensland Pittosporum
P. tobira	Japanese Pittosporum
Podocarpus macrophyllus	Japanese Yew
Prunus laurocerasus	English Laurel
Prunus spp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
Pseudotsuga menziesii	Douglas Fir**
Pyracantha coccinea	Firethorn (Scarlet)
P. fortuneana	Firethorn
P. koidzumii	Firethorn
Pyrus spp.	Bradford Pear spp.
Quercus Shumardii	Oak, Shumard's Red
Quercus spp.	Oak species
Raphiolepis indica	Indian Hawthorne
Raphiolepis umbellata	Yedda Hawthorne
Rhododendron (including Azalea)	'Coral Bells,' 'Formosa,' 'Hino-crimson,' 'PJM' 'Delaware Valley White,' 'Roseum Elegans,' 'Flame Creeper,' 'Girard Crimson,' 'Wakeiebisu' 'White Gumbo,' 'George L. Tabor'
Rosa banksiae	Lady Bank's Rose
Rosmarinus officinalis	Rosemary*
Rudbeckia spp.	Black-Eyed Susan; Goldsturm
Santolina virens	Santolina
Saxifraga spp.	Saxifrage; Purple Dome
Scabiosa spp.	Pincushion Flower
Sedum album	Stonecrop
Sedum caudicicola	Stonecrop; Lidakense
Sedum dasyphyllum	Stonecrop
Sedum spurium	Stonecrop; Dragon's Blood
Spiraea bumalda	Spirea; Anthony Waterer
Syzygium paniculatum	Australian Brushcherry
Taxus cuspidata	Japanese Yew
T. media	Yew
Teucrium spp.	Germander
Thalictrum dipterocarpum	Meadow Rue
Thuja occidentalis	American Arborvitae
Trachelospermum asiaticum	Star Jasmine
Tsuga canadensis	Canada Hemlock
Tulipa spp.	Tulip species
Veronica spp.	Veronica, Speedwell; Sunny Border
Viburnum japonicum	Japanese Viburnum
V. odoratissimum	Sweet Viburnum
V. plicatum	Japanese Snowball
V. rigidum	Canary Island Viburnum
V. suspensum	Arrowood Viburnum
V. tinus	Laurustinus
V. trilobium	Cranberry Bush
V. wrightii	Leatherleaf Viburnum
Vinca major	Greater Periwinkle, Vinca
Vinca Minor	Common Periwinkle, Vinca
Vitis spp.	Grape*
Weigela florida	Old Fashioned Weigela
Yucca aloifolia	Spanish Bayonet, Yucca

Y. filametosa

Adam's Needle, Yucca

\*Do not use on food producing trees, vines or plants.

\*\*Not for use on container grown plants.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container in a dry place. Store in locked area out of reach of children and domestic animals.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

**CONTAINER Disposal:** Non-refillable container. Do not reuse or refill this container. Empty container through use according to label instructions. Do not rinse unless required for recycling. Offer for recycling, if available. If recycling is not available; then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

#### CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop, injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of Control Solutions, Inc. or the Seller. All such risks shall be assumed by the Buyer.

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label, and is reasonably fit for the purposes set forth in the DIRECTIONS FOR USE, when it is used in accordance with such directions, subject to the inherent risks mentioned above.

CONTROL SOLUTIONS, INC. NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THIS WARRANTY EXTENDS TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNING, OR CAUTIONS. BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. CONTROL SOLUTIONS, INC. and the seller offer this product, and the Buyer and User accept it, subject to the foregoing CONDITIONS OF SALE AND WARRANTY.

#### *Optional State Specific Language*

[This product may stain driveways, walks, patios, etc.]

[Wash or sweep off thoroughly, immediately after application]

[Do Not Apply Near Water, Storm Drains or Drainage Ditches.]

[Do Not Apply if heavy rain is expected]

[Apply this product only to your lawn or garden, and sweep any product that lands on the driveway, sidewalk or street, back onto your lawn or garden]

[Help do your part in protecting our rivers and lakes by following label directives. Do not apply near water storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk or street back onto your lawn. Do not apply to frozen ground.]

{Label may also include state specific language regarding Nitrogen application}